

Spinal cord compression associated with pseudohypoparathyroidism

S M Alam MB BS MRCP **W Kelly** MD MRCP
*Department of Endocrinology and Diabetes,
 Middlesbrough General Hospital,
 Ayresome Green Lane, Middlesbrough TS5 5AZ*

Keywords: spinal cord compression; pseudohypoparathyroidism; epilepsy

Pseudohypoparathyroidism was described by Albright in 1942¹. The features include obesity, mental retardation, epilepsy, and hypocalcaemia. Resistance to parathyroid hormone (PTH) was suggested¹, and later verified by PTH infusions. We report a rare neurological association.

Case report

The weight, height and bone age of the patient were above average at 3½ years. At 11 years the patient had epilepsy, controlled by phenytoin. At 23 years he had four weeks of progressive leg weakness, then incontinence of urine and faeces. His appearance suggested Cushing's syndrome, with facial roundness, obesity, weakness, and abdominal striae (Figure 1). He was hypogonadal with undescended testes. He had a spastic paraparesis without a sensory level.

Serum calcium was low, and alkaline phosphatase was raised (Table 1). Plasma bioactive PTH was low, and when 100 ng/l of PTH standard was added to plasma samples from the patient, only 35% was recovered (normal, 50–90%), suggesting an inhibitor to the action of PTH². Bovine PTH intravenously showed impaired phosphate excretion and a sub-normal rise in urinary cAMP. With alpha calcidol therapy, serum calcium became normal. The myelogram showed compression at T1/2, T10/11 and L4/5. Bone histology showed no features of osteomalacia or hyperparathyroidism.

The spinal cord was decompressed by three separate dorsal laminectomies. Very hard bone was noted, but no evidence of extradural or intradural space-occupying lesions. The cord looked normal. There was initially rapid improvement in

Table 1. Investigations in a patient with pseudohypoparathyroidism

	Inpatient	Normal range
Serum calcium (mmol/l)	1.38	2.1–2.6
Serum albumin (g/l)	45	35–50
Serum inorganic phosphate (mmol/l)	1.75	0.8–1.4
Alkaline phosphatase (u/l)	142	35–115
Plasma immunoactive PTH levels (mean=0.6, SD=0.3, n=6) (u/l)	0.3 & 0.9	0.3–1.4
Plasma bioactive PTH concentration (1 ng/l=2.3×10 ⁻³ IU/l) (ng/l)	0.15	1–15
Urinary calcium excretion (mmol/24 h)	0.34–0.83	2.5–7.5
Phosphate excretion (mmol/24 h)	29	16–50
Serum 25-hydroxyvitamin D (pre-treatment), September 1986 (nmol/l)	7	25–75
Testosterone (nmol/l)	3.1	11–36
Urea and electrolytes	normal	
Urinary free cortisol (nmol/24 h)	220	<300
Plasma cortisol 0900 h (nmol/l)	140	150–600
Free thyroxine index (units)	104	55–180

bladder and bowel function, and a gradual improvement in the function of the legs. With physiotherapy the patient could walk with a frame. He then deteriorated, with weaker legs. Further neurosurgical decompression was attempted, but the spinal cord had infarcted below T10, and he is again paraplegic.

Discussion

Our patient presented with the typical features of pseudohypoparathyroidism type 1¹. Treatment with alpha calcidol increased serum calcium, and decreased alkaline phosphatase.

Patients with pseudohypoparathyroidism have been reported to have a spectrum of radiological abnormalities from osteoporosis to osteitis fibrosa cystica and osteomalacia³. Sometimes the radiological appearance is normal, but osteomalacia has been suspected on biochemical evidence, and confirmed by bone histology⁴. The normal bone histology and the apparently normal or low immunoactive PTH in our patient may be due to the presence of the PTH inhibitor². There have only been four previous reports of patients with spinal cord compression and pseudohypoparathyroidism^{5–8}. These authors reported excessive bone formation in the vertebral canal at C3/4⁷ and spondylogenic myelopathy leading to paraplegia⁶.

The aetiology of cord compression in pseudohypoparathyroidism is disputed. Calcification of the longitudinal ligaments of the spine may compress the cord⁸. The cord compression may be related to the effects of PTH and vitamin D on the vertebrae and ligaments. It is interesting that X-linked hypophosphataemic rickets may also be complicated by spinal cord compression and extremely hard bone⁹, both of which occur in pseudohypoparathyroidism, as in our case⁵. Although he presented with low 25-OH-vitamin D, our patient had no radiological or histological evidence of osteomalacia.

It has also been suggested that the cord compression develops as a result of vitamin D therapy⁵; this seems unlikely, since patients have developed the complication before vitamin D has been given, and myelopathy has not worsened after vitamin D therapy has been commenced. Possibly the bony abnormalities of the spine are genetically determined. The response to surgical decompression of the cord has been disappointing. Therefore, clinicians who follow up these patients should be aware of this complication, and early diagnosis is essential.

Acknowledgments: Biological activity of PTH and the inhibitor were assessed by Dr Loveridge, now at Rowett Research Institute, Aberdeen. We thank our colleagues in the Departments of Neurology, Neurosurgery, Radiology and Biochemistry.

Case presented to Section of Endocrinology, 27 May 1987




Figure 1. Patient aged 23 years, showing obesity (141 kg) and striae. This classical appearance of pseudohypoparathyroidism resembles Cushing's syndrome, but cortisol levels were not elevated

0141-0768/90/
 010060-02/\$02.00/0
 © 1990
 The Royal
 Society of
 Medicine

References

- 1 Albright F, Burnett CH, Smith PH, Parson W. Pseudohypoparathyroidism: An example of 'Seabright-Bantam Syndrome'. *Endocrinology* 1942;30:922-32
- 2 Loveridge N, Bradbeer J, Dunham C, de Deuxchaisnes CN, Fischer JA. Pseudohypoparathyroidism type I: disparity between bio-PTH levels measured by the renal and metatarsal cytochemical bioassays. In: Cohn DV, Martin TJ, Meunier PJ, eds. *Calcium regulation and bone metabolism: Basic and clinical aspects*, 1987;9:133-7
- 3 Burnstein MI, Kottamasu SR, Pettifor JM, Sochett E, Ellis BI, Frame B. Metabolic bone disease in pseudohypoparathyroidism: radiologic features. *Radiology* 1985; 155:351-6
- 4 Epstein S, Meunier PJ, Lambert PW, Stern PH, Bell NH. 1-alpha-25-dihydroxyvitamin D3 corrects osteomalacia in hypoparathyroidism and pseudohypoparathyroidism. *Acta Endocrinol (Copenh)* 1983;103:241-7
- 5 Cullen DR, Pearce JM. Spinal cord compression in pseudohypoparathyroidism. *J Neurol Neurosurg Psychiatry* 1964;27:459-62
- 6 Greco R, Sagratella S, Landucci C, Canova A. Paraplegia da mielopatia spondilogenica in soggetto con osteodistrofia di Albright tipo II o pseudo-pseudoiparathyroidismo. *Riv Neurobiol* 1978; 24:243-54 (summary in English).
- 7 Cavallo A, Meyer WJ, Bodensteiner JB, Chesson AL. Spinal cord compression: an unusual manifestation of pseudohypoparathyroidism. *Am J Dis Child* 1980;134:706-7
- 8 Firooznia H, Golimbu C, Raffi M. Case report 312. Diagnosis: progressive paraparesis in a woman with pseudohypoparathyroidism with ossification of the posterior longitudinal ligament from C4 to T5. *Skeletal Radiol* 1985;13:310-13
- 9 Bradbury PG, Brenton DP, Stern GM. Neurological involvement in X-linked hypophosphataemic rickets. *J Neurol Neurosurg Psychiatry* 1987;50:810-12

(Accepted 29 March 1989. Correspondence to Dr W F Kelly)

Primary intraocular lymphoma with lymphomatous meningitis

P G Hykin MBBS SHO J E MacSweeney MRCP
D Gartry FRCS E Graham MRCP DO *Medical
Eye Unit, St Thomas' Hospital, London SE1 7EH*

Keywords: intraocular lymphoma; lymphomatous meningitis; seventh nerve palsy

We report a patient with ocular lymphoma who developed bilateral seventh nerve palsies due to lymphomatous meningitis without intracerebral involvement. The palsies markedly improved after cerebral irradiation and intrathecal methotrexate. Further studies are required to determine whether repeated cerebrospinal fluid examination and prophylactic cranial and spinal radiotherapy are indicated in patients who present with the disease localized to the eye.

Case report

A 62-year-old man had blurred vision for 2 years, diagnosed as bilateral posterior uveitis. Aetiological investigations were unrewarding. Despite repeated courses of high dose oral prednisolone, 20 months later visual acuity and colour vision deteriorated and he suddenly developed bilateral seventh nerve palsies.

At presentation, right visual acuity was 6/9, left 6/24. No Ishihara colour plates were seen with either eye. Pupillary responses were normal and there was no relative afferent pupillary defect. Both visual fields were markedly constricted. The anterior chambers were quiet. There were numerous large cells in both vitreous chambers, with a resultant hazy fundal appearance. Fluorescein angiography showed characteristic pigment epithelial defects. There were marked bilateral seventh nerve palsies (Figure 1) with left hyperacusis. There was no clinical evidence of lymphoma elsewhere.

Cytocentrifuge and immunochemical staining of cerebrospinal fluid revealed β -cell lymphoma cells (Figure 2). Chest X-ray, bone marrow trephine, and computerized axial tomography of the head and abdomen were normal.




Figure 1. Bilateral seventh nerve palsies

Orbital irradiation (600 cGy) and total cranial irradiation (3750 cGy) were given. After therapy, right visual acuity was 6/6, left 6/18. Both vitreous chambers had cleared. Both facial palsies were markedly improved and the patient remained well for 2 years before progressive dementia and death from subsequent cerebral lymphoma.

Discussion

This patient presented with posterior uveitis of obscure aetiology. Bilateral uveitis, failure to respond to steroids and the high incidence of ocular lymphoma in this age group were suggestive of the diagnosis¹.

Case presented to
Section of
Ophthalmology,
12 May 1988

0141-0768/90/
010051-02/\$02.00/0
© 1990
The Royal
Society of
Medicine